

Sources affecting knowledge and behavior responses to the Zika virus in US households with current pregnancy, intended pregnancy and a high probability of unintended pregnancy

Man-pui Sally Chan¹, Mohsen Farhadloo^{1,2}, Kenneth Winneg², Kathleen Hall Jamieson², Dolores Albarracin¹

¹Department of Psychology, University of Illinois at Urbana-Champaign, 603 E. Daniel St, Champaign, IL 61820, USA

²Annenberg Public Policy Center, University of Pennsylvania, 202 S 36th St, Philadelphia, PA 19104, USA

Address correspondence to Man-pui Sally Chan, E-mail: sallycmp@illinois.edu

ABSTRACT

Background This study examined the influences of information sources on Zika-relevant knowledge and behaviors in US households containing members who are pregnant, intend to become pregnant, or have a higher probability of unintended pregnancy in Zika-affected regions (i.e. respondents who are younger, are black, have less education, are unmarried, and reside in the southern USA).

Methods Over 32 000 US adults completed a survey measuring Zika-relevant knowledge and behaviors along with information sources (e.g. discussing Zika with practitioners), general media usage (e.g. TV) and demographic information over 30 weeks.

Results Respondents in the group with (versus without) either pregnancy or intended pregnancy were more likely to use all information sources, which in turn created differences in knowledge and behavior responses. To gauge information sources in US-South respondents with a high probability of having a household member with unintended pregnancy based on demographics, younger, less-educated, unmarried, black respondents had fewer Zika discussion with practitioners than another group.

Conclusions Efforts to increase Zika-related knowledge and protective behaviors should target households with members who are pregnant or intending to become pregnant via practitioners, family and friends. Additional efforts should target information channels to reach younger, less educated, unmarried, black respondents, which are at risk for unintended pregnancy.

Keywords attitude, behavior, communication, health, information sources, knowledge, pregnancy, public health, Zika

Introduction

The Zika virus is a mosquito-borne flavivirus recently related to a number of human diseases, including microcephaly and possibly Guillain-Barre syndrome.¹ On 1 February 2016, the World Health Organization declared the ongoing Zika crisis an emergency.² The virus is primarily transmitted by the bite of the female *Aedes* mosquito species, the same vector that transmits yellow fever, dengue fever and chikungunya. Zika can be passed from a mother to her fetus during pregnancy and can also be sexually transmitted from infected people to their partners. In 8 out of 10 cases, the person carrying the virus is asymptomatic. Infection with Zika is diagnosed based on a blood or urine test. There is no available treatment to cure or vaccine to prevent infections, which increases the

need to understand how to best inform the population on transmission and preventive practices.^{2–4} As of 18 January 2017, the Zika virus infections are notifiable conditions in the USA, which has resulted in 4900 reported cases, 96% associated with traveling outside the USA.^{5,6}

Even though infection during pregnancy can result in several fetal impairments (i.e. microcephaly, defects of eyes, hearing deficits and impaired growth).^{7,8} US studies have suggested low levels of knowledge about Zika transmission

Man-pui Sally Chan, Research Assistant Professor of Psychology

Mohsen Farhadloo, Postdoctoral Research Associate

Kenneth Winneg, Managing Director of Survey Research

Kathleen Hall Jamieson, Elizabeth Ware Packard Professor of Communication

Dolores Albarracin, Professor of Psychology, Business, and Medicine

and limited practice of behaviors that help to prevent infections.^{9–11} Therefore, increasing population knowledge about Zika and about means to prevent Zika infection is critical from a public health perspective.^{4,12} Key populations involve members of households with pregnant women, women who intend to get pregnant, and women who might unintentionally become pregnant based on respondents' demographic characteristics and who live in the South of the USA (i.e. an area of active Zika virus transmission).^{5,13} This article presents the first survey continuously administered by phone (landlines and cell phones) to a nationally representative sample in the USA over 30 weeks from 16 February to 4 September in 2016.

The present study contributes to previous work by (i) examining different measures of knowledge, behavioral intentions and behaviors, such as changing travel plans and wearing mosquito repellent to reduce the chance of mosquito-bites, and (ii) determining the likely sources of knowledge and behavioral responses, including discussions with practitioners, discussions with friends and family, seeking information online, and general media use (e.g. social media, newspapers, online news, television and radio) in samples with or without either pregnancy or intended pregnancy (i.e. samples from households with either pregnancy or intended pregnancy versus neither pregnancy nor intended pregnancy), samples from households with pregnancy intentions (i.e. the pregnancy-intention- versus current-pregnancy-groups), and samples with demographic characteristics associated with unintended pregnancy (i.e. less-educated unmarried black respondents versus other respondents) who also live in Zika-affected regions. Importantly, unintended pregnancy rates are highest among unmarried women, women who are black, and women with less education/income.^{14–17} Therefore, we selected younger respondents who reside in the region of active Zika transmissions and have household members with neither current nor intended pregnancy and then compared those who are less-educated, unmarried, and black with others to understand the information sources of this at-risk population as well. The guiding research questions of our analyses included: Are there differences in information sources across groups? Do these information sources mediate differences in knowledge and behaviors across these samples?

Method

Sample and survey design

We conducted a survey of knowledge, behavioral intentions and behaviors relevant to Zika over 30 weeks. Each weekly,

dual-frame sample was designed to represent the adult US population (including Hawaii and Alaska) and used a fully-replicated, single-stage, random-digit-dialing (RDD) sample of landline telephone households, along with randomly generated cell phone numbers. Each weekly wave consisted of 1000 interviews of which at least 600 were obtained from cell phone respondents. Within each landline household, a single respondent (youngest adult) was selected. Cell-phone respondents were considered separately from landlines as the interview may take place outside the respondent's home. Surveys were conducted over a 5-day period, in English and Spanish, typically from Wednesday through Sunday, to include both weekdays and weekends. Each weekly wave was weighted to provide nationally representative and projectable estimates of the adult population 18 years of age and older. The weighting process takes into account the disproportionate probabilities of household and respondent selection due to the number of separate telephone landlines and cell phones answered by respondents and their households, as well as the probability associated with the random selection of an individual household member. Following application of the above weights, the sample is post-stratified and balanced by key demographics such as age, race/ethnicity, sex, the region of residence and education. The sample is also weighted to reflect the distribution of phone usage in the general population, meaning the proportion of those who are cell phone only, landline only and mixed users. The average response rate over the weeks was 7.5%, and over 32 800 US adults completed the survey over 30 weeks. (More details about the survey design can be found in the Supplementary material).

We combined respondents' answers to questions about the pregnancy status and intentions to get pregnant by members in the households, together with questions about demographic characteristics (i.e. age, the highest education level attained, race/ethnicity, and marital status), and the geographic residence region to identify vulnerable samples. The first sample contained respondents from households with versus without members who are either pregnant or intending to become pregnant (i.e. samples from households with either pregnancy or intended pregnancy versus neither pregnancy nor intended pregnancy), $N = 24\,459$. The second sample consists of respondents from households with members, who are currently versus intending to get pregnant (i.e. the pregnancy-intention- versus current-pregnancy-groups), $N = 1465$. The third sample includes younger respondents (i.e. aged below 25), who live in Zika-affected regions (i.e. the Southern regions) and have household members, who do not intend to become pregnant. Specifically, the third sample includes less-educated (i.e. less than a college degree)

Table 1 Descriptive statistics of all samples

Variable	First		Second		Third	
	Either current or intended pregnancy	Neither	Pregnancy-intention-group	Current-pregnancy-group	Less-educated unmarried black respondents	Other respondents
<i>N</i>	1568	22 891	947	518	122	758
Sex	50% females	51% females	51% females	50% females	46% females	41% females
Age						
Aged 24 or below	11%	10%	10%	12%	100%	100%
Aged 25 or above	89%	90%	90%	88%	0%	0%
Education						
Without a college degree	45%	40%	50%	38%	100%	77%
With a college degree or a postgraduate/professional degree	55%	60%	50%	62%	0%	23%
Ethnicity						
White	69%	76%	71%	68%	0%	66%
Black	11%	11%	10%	12%	100%	5%
Hispanics	20%	12%	18%	21%	12%	29%
At-risk region	42%	43%	43%	42%	100%	100%
Media usage						
Forwarding information online	19%	11%	18%	21%	10%	15%
Reading newspaper online and offline	3.99 (2.07)	4.05 (2.06)	4.05 (2.05)	3.91 (2.11)	3.36 (2.08)	3.36 (1.92)
Using online news sites	3.15 (2.23)	2.65 (2.11)	3.24 (2.24)	2.96 (2.20)	2.62 (1.93)	3.00 (2.07)
Listening to radio online and offline	4.15 (2.19)	3.89 (2.22)	4.20 (2.17)	4.04 (2.22)	3.74 (2.18)	3.73 (2.18)
Watching television online and offline	5.06 (1.90)	5.30 (1.79)	5.11 (1.87)	5.06 (1.88)	4.81 (2.01)	4.74 (1.93)
Usage frequency of Twitter	1.81 (1.68)	1.59 (1.46)	1.84 (1.70)	1.74 (1.62)	2.92 (2.35)	2.55 (2.18)
Usage frequency of Facebook	4.52 (2.35)	3.79 (2.45)	4.60 (2.33)	4.41 (2.38)	4.49 (2.20)	4.98 (2.10)
Seeking information about Zika online	46%	26%	47%	43%	33%	42%
Discussing Zika with practitioners	18%	7%	16%	22%	2%	10%
Discussing Zika with family and friends	43%	32%	42%	45%	26%	29%
Knowledge						
Sexually transmitted	2.99 (1.10)	2.84 (1.10)	2.97 (1.10)	2.99 (1.10)	2.82 (1.09)	2.89 (1.04)
Mosquito-transmitted	3.37 (0.86)	3.31 (0.87)	3.37 (0.85)	3.36 (0.87)	3.44 (0.79)	3.43 (0.76)
Microcephaly	3.40 (0.81)	3.35 (0.79)	3.42 (0.79)	3.37 (0.82)	2.97 (0.94)	3.14 (0.88)
Noticeable symptoms	2.67 (0.97)	2.67 (0.95)	2.72 (0.96)	2.63 (0.96)	2.37 (0.96)	2.58 (0.89)
Change of travel plans	3.22 (1.09)	3.05 (1.13)	3.14 (1.12)	3.37 (1.00)	3.42 (0.84)	3.10 (1.02)
Get vaccinated if available	2.80 (1.16)	2.50 (1.17)	2.80 (1.16)	2.84 (1.15)	2.80 (1.07)	2.84 (1.00)
Taking protective behaviors	35%	26%	32%	40%	21%	27%

unmarried black respondents, $N = 880$. Table 1 presents descriptive characteristics of different samples. Details about the survey items appear next.

Measures

Demographics

Respondents answered questions about their sex, age, marital status, the highest education level they attained, and their region of residence. We then categorized their responses into groups: age = aged 24 or below versus aged 25 or above; marital status = unmarried (i.e. single never married or cohabiting) versus married, education = Without a college degree versus with a college degree or a postgraduate/professional degree; and region of residence = US's south where reported most Zika infections versus other US regions.

Household with current or intended pregnancy

We used two questions with a dichotomous response of yes and no to determine the pregnancy grouping of the respondents' household, i.e. as far as you know, is anyone in your household currently pregnant?, and, as far as you know, is anyone in your household considering getting pregnant within the next 12 months?, in different samples. The sample of pregnancy or intended pregnancy contains respondents who answered yes to both or either of the questions (coded as -1) and is contrasted with respondents who answered no to both questions (coded as 1). The pregnancy-intention-group includes those who answered yes to the second question (coded as 1) and is contrasted with the current-pregnancy-group (coded as -1), which includes respondents who answered yes to the first question. The unintended-pregnancy-probability-sample includes younger respondents who reside in the South of the USA and answered no to both questions. This sample contrasts respondents who are less-educated, unmarried and black (coded as 1) with those without these characteristics (coded as -1).

Sources of Zika information and media usage

Respondents also reported whether they discussed Zika with practitioners, discussed it with family and friends, sought such information online, and used other media, including forwarding information about Zika online, using online news sites, reading newspapers online or offline, listening to public radio online or offline, watching television online or offline, and using social media such as Twitter and Facebook. The question about forwarding information about Zika online had the dichotomous choices of yes and no, whereas the

other questions had a polychotomous scale (1 = never to 7 = many times a day) on which to indicate use.

Knowledge

Respondents answered four questions on a polychotomous scale (1 = not at all accurate/likely to 4 = very accurate/likely), to indicate their level of knowledge of Zika (see Table SI). Two questions were about how Zika is transmitted: Mosquito bites and sex. The other questions were about Zika infection symptoms and outcomes, including the association between Zika and microcephaly and whether or not Zika infections always produce noticeable symptoms (this is an incorrect item, and the responses were reverse coded to indicate the level of accurate knowledge).

Behavioral intentions and behaviors

For behaviors related to Zika, respondents answered questions about the likelihood that they would change their travel plans if they learned about an outbreak of Zika in their travel destination, and the likelihood that they would receive a Zika vaccine if one were available (see Table SI). Furthermore, respondents were asked an open-ended question about whether they engaged in any behaviors to protect them from becoming infected with Zika in the past 3 months. The protective behaviors listed included purchasing/wearing insect repellent, wearing long-sleeved shirts and pants, and removing standing water.

Statistical analyses

The number of weeks for which each question was asked in the survey was not identical because some questions were rotated in and out of the survey during the survey period. Given the presence of systematic missing data, it is inappropriate to apply any imputation techniques, which assume the missing data to be random. Therefore, we used ordinal logistic regression to analyze each knowledge and behavior response individually.

There were two steps in the statistical analyses: We first conducted logistic regressions to examine the relations between critical samples of comparison and sources of Zika information (i.e. path *a*), and then carried out ordinal logistic regressions to delineate the relations of sources of Zika information, including seeking information about Zika online, discussing Zika with practitioners, and discussing Zika with family and friends, with knowledge and behavior outcomes, while controlling for the indicators for the critical samples (i.e. path *b*). Ordinal logistic regression is an extension of logistic regression when the outcome is ordinal and has more than two responses. Therefore, we used logistic regression to analyze

the behavior response—taking protective behaviors, due to its dichotomous nature. The same set of analyses was repeated separately for the three comparisons to make determinations about vulnerable groups (see Appendix A for additional comparisons of vulnerable groups with specific sociodemographic characteristics).

For the first analysis, which compared groups with or without either current or intended pregnancy, we examined whether the group variable was associated with information sources, and in turn, whether these sources led to differential knowledge and behavior outcomes between the groups. Rather than treating current and intended pregnancy as part of the same group, the second analysis compared the pregnancy-intention- versus current-pregnancy-groups. The third analysis focused on the unintended-pregnancy-probability-sample of younger respondents at Zika-infected regions. This analysis included two groups (i.e. less-educated unmarried black versus other respondents), which were entered in the analyses to study the relations with information sources of Zika and their mediating roles on knowledge and behavior outcomes.

Additionally, we included the main effects of sex, the week number during which the survey was completed, and media usage (i.e. forwarding information about Zika online, using online news sites, reading newspapers online or offline, listening to public radio online or offline, watching television online or offline, and using social media such as Twitter and Facebook), as well as the interaction effects of information sources with sex, of information sources with the group variable, and of information sources with the region of residence (except the third sample) as covariates in each analysis.

For all three analyses, when paths *a* and *b* were both significant, we performed a Sobel test to determine whether the included mediator (i.e. seeking information about Zika online, discussing Zika with practitioners, or discussing Zika with family and friends) was significant. The Sobel test computes the two regression paths described above (i.e. paths *a* and *b*), together with the variances as a ratio, and treats the ratio as a χ -test. These analyses test whether an information source mediates differences in knowledge of behavior between those groups.

Results

Tables 2–4 present the results of three critical samples. Path *a* indicates whether there were any between-group differences in information sources. Path *b* represents the correlation of information sources with knowledge and behavior responses. Path *c* shows the association of the group

variable with knowledge and behavior responses when taking the mediator into account. The analyses of each critical sample had different *N*s because of different sampling criteria and the presence of missing responses to different survey items. The *N*s of each knowledge and behavior responses ranged: Sexually transmitted = 872–14 571, mosquito-transmitted = 904–15 452, microcephaly = 932–15 970, noticeable symptoms = 924–15 822, change of travel plans = 709–12 274, get vaccinated if available = 361–5 918, and taking protective behaviors = 656–11 544.

Are there differences in information sources as a function of the critical samples? Do information sources mediate group differences in knowledge and behaviors?

The comparison of samples from households with either pregnancy or intended pregnancy versus neither pregnancy nor intended pregnancy

Table 2 shows significant differences in all information sources between groups with or without either pregnancy or intended pregnancy (i.e. path *a*, *P*s < 0.001). Furthermore, some information sources correlated with knowledge and behaviors (path *b*). For example, seeking Zika information online and discussing Zika with family and friends were associated with knowledge about sexual transmission and microcephaly, and with behavioral intentions to change travel plans. Additionally, several of the knowledge and behavior responses (i.e. sexually transmitted, microcephaly, change of travel plans, get vaccinated if available and taking protective behaviors) differed across groups as well (see Table SII), thus comprising plausible mediations of the differences in knowledge and behavior.

As revealed by the Sobel tests, all three information sources significantly explained the differences between groups with or without either pregnancy or intended pregnancy in knowledge and behavior outcomes (Table 2). For example, seeking information online completely mediated (i.e. path *c* was non-significant) the link between the group variable and knowledge about microcephaly, Sobel test = -2.08 , *P* = 0.0375, and changing travel plans, Sobel test = -2.92 , *P* = 0.0035. Moreover, discussing Zika with practitioners completely mediated the links between the group variable and two knowledge outcomes, i.e. the mosquito-transmitted nature of Zika, Sobel test = -2.20 , *P* = 0.0278 and not always having noticeable symptoms, Sobel test = -2.27 , *P* = 0.0232, while discussing it with family and friends only showed a complete mediation of group differences in knowledge of microcephaly, Sobel test = -2.07 , *P* = 0.0385. In summary, the group with (versus without) either pregnancy or intended pregnancy was more likely to seek information

Table 2 Results of mediation analyses of the comparison between samples from households with either current or intended pregnancy and samples from households with neither current nor intended pregnancy

<i>Mediator</i>	<i>Knowledge</i>															
	<i>Sexually transmitted</i>				<i>Mosquito-transmitted</i>				<i>Microcephaly</i>				<i>Noticeable symptoms</i>			
	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>
Seeking Zika information online	-0.44 (0.08)***	0.31 (0.13)*	-0.19 (0.09)*	-2.19*	-0.42 (0.08)***	0.09 (0.13)	-0.11 (0.09)	-	-0.50 (0.07)***	0.30 (0.14)*	-0.16 (0.09)	-2.08*	-0.48 (0.07)***	0.20 (0.12)	-0.00 (0.08)	-
Discussing Zika with practitioners	-0.79 (0.09)***	0.14 (0.17)	-0.22 (0.07)***	-	-0.78 (0.09)***	-0.39 (0.17)*	0.13 (0.07)	2.20*	-0.75 (0.09)***	0.20 (0.18)	-0.13 (0.07)	-	-0.73 (0.09)***	0.38 (0.16)*	0.02 (0.07)	-2.27*
Discussing Zika with family and friends	-0.32 (0.07)***	0.31 (0.13)*	-0.17 (0.09)*	-2.09*	-0.28 (0.07)***	0.11 (0.13)	-0.11 (0.09)	-	-0.33 (0.07)***	0.31 (0.13)*	-0.15 (0.09)	-2.07*	-0.33 (0.07)***	0.12 (0.12)	0.09 (0.08)	-

<i>Mediator</i>	<i>Behavioral intentions and behaviors</i>											
	<i>Change travel plans</i>				<i>Get vaccinated if available</i>				<i>Taking protective behavior(s)</i>			
	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>	<i>Path a</i>	<i>Path b</i>	<i>Path c'</i>	<i>Sobel test</i>
Seeking Zika information online	-0.35 (0.09)***	0.67 (0.16)***	-0.12 (0.10)	-2.92***	-0.54 (0.12)***	0.21 (0.20)	-0.33 (0.13)**	-	-0.36 (0.09)***	0.43 (0.18)*	-0.33 (0.12)***	-2.01*
Discussing Zika with practitioners	-0.69 (0.10)***	0.10 (0.20)	-0.3 (0.08)***	-	-0.85 (0.14)***	0.51 (0.26)*	-0.30 (0.11)**	-1.86	-0.66 (0.11)***	1.00 (0.23)***	-0.27 (0.10)**	-3.51***
Discussing Zika with family and friends	-0.36 (0.08)***	0.34 (0.15)*	-0.31 (0.10)***	-2.00*	-0.34 (0.12)***	0.44 (0.20)*	-0.26 (0.13)*	-1.76	-0.38 (0.08)***	0.68 (0.18)	-0.26 (0.13)*	-2.86***

Note. Samples from households with neither current nor intended pregnancy were coded as 1 whereas samples from households with either current or intended pregnancy were coded as -1. Unstandardized estimates and standard errors in parentheses. Main effects of sex, age group, level of education, at-risk region, week number of the survey completion, ethnicity, seeking Zika information online, discussing Zika with practitioners, discussing Zika with family and friends, forwarding information online, reading newspapers online and offline, using online news sites, listening to radio online and offline, watching television online and offline, usage frequency of Twitter, and usage frequency of Facebook, were included for the estimations of paths *a* and *b*, while the interaction effects between the mediators and the group variable, sex and at-risk region were included for the estimation of paths *b* and *c'*.

*** $P \leq 0.001$. ** $P \leq 0.01$. * $P \leq 0.05$.

Table 3 Results of mediation analyses of the comparison of the pregnancy-intention- versus current-pregnancy-groups

Mediator	Knowledge															
	Sexually transmitted				Mosquito-transmitted				Microcephaly				Noticeable symptoms			
	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test
Seeking Zika information online	0.31 (0.17)	0.12 (0.28)	-0.28 (0.18)	-	0.26 (0.16)	0.43 (0.3)	0.18 (0.18)	-	0.29 (0.16)	-0.20 (0.30)	-0.08 (0.19)	-	0.26 (0.16)	-0.04 (0.27)	0.08 (0.17)	-
Discussing Zika with practitioners	-0.52 (0.19)**	0.35 (0.34)	-0.05 (0.15)	-	-0.55 (0.18)***	-0.16 (0.36)	0.15 (0.16)	-	-0.47 (0.18)**	-0.27 (0.36)	-0.12 (0.16)	-	-0.48 (0.18)**	-0.21 (0.33)	0.08 (0.14)	-
Discussing Zika with family and friends	-0.26 (0.16)	0.49 (0.28)	-0.10 (0.18)	-	-0.26 (0.15)	0.41 (0.29)	0.27 (0.19)	-	-0.26 (0.15)	0.41 (0.29)	0.15 (0.19)	-	-0.25 (0.15)	-0.24 (0.26)	-0.03 (0.18)	-
Mediator	Behavioral intentions and behaviors															
	Change travel plans				Get vaccinated if available				Taking protective behavior(s)							
	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test				
Seeking Zika information online	0.38 (0.19)*	0.25 (0.36)	-0.64 (0.21)***	-	0.16 (0.26)	0.65 (0.45)	0.23 (0.29)	-	0.21 (0.20)	0.51 (0.42)	-0.19 (0.25)	0.31				
Discussing Zika with practitioners	-0.44 (0.22)*	-0.02 (0.46)	-0.47 (0.19)**	-	-0.03 (0.30)	0.82 (0.55)	0.35 (0.24)	-	-0.40 (0.23)	1.27 (0.52)**	-0.38 (0.22)	-1.47				
Discussing Zika with family and friends	-0.42 (0.18)*	0.32 (0.35)	-0.35 (0.23)	-	-0.16 (0.25)	1.23 (0.44)	0.50 (0.29)	-	-0.32 (0.19)	0.90 (0.43)*	-0.2 (0.29)	-1.25				

Note. The pregnancy-intention-samples were coded as 1 whereas the current-pregnancy-samples were coded as -1. Unstandardized estimates and standard errors in parentheses. Main effects of sex, age group, level of education, at-risk region, week number of the survey completion, ethnicity, seeking Zika information online, discussing Zika with practitioners, discussing Zika with family and friends, forwarding information online, reading newspapers online and offline, using online news sites, listening to radio online and offline, watching television online and offline, usage frequency of Twitter, and usage frequency of Facebook, were included for the estimations of paths a and b, while the interaction effects between the mediators and the group variable, sex and at-risk region, were included for the estimation of paths b and c'.

*** $P \leq 0.001$. ** $P \leq 0.01$. * $P \leq 0.05$.

Table 4 Results of mediation analyses of the unintended-pregnancy-probability-sample of younger respondents at Zika-infected regions

Mediator	Knowledge															
	Sexually transmitted				Mosquito-transmitted				Microcephaly				Without noticeable symptoms			
	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test
Seeking Zika information online	-0.46 (0.29)	0.11 (0.22)	0.05 (0.28)	-	-0.48 (0.29)	0.15 (0.24)	0.57 (0.32)	-	-0.5 (0.28)	0.15 (0.23)	-0.21 (0.27)	-	-0.5 (0.28)	0.15 (0.23)	-0.21 (0.27)	-
Discussing Zika with practitioners	-2.27 (1.03)*	0.56 (0.35)	0.08 (0.23)	-	-2.34 (1.03)*	0.65 (0.39)	0.28 (0.25)	-	-1.72 (0.74)*	-0.06 (0.35)	-0.2 (0.23)	-	-1.72 (0.74)*	-0.06 (0.35)	-0.2 (0.23)	-
Discussing Zika with family and friends	-0.27 (0.29)	0.42 (0.23)	-0.01 (0.28)	-	-0.23 (0.28)	0.38 (0.25)	0.35 (0.3)	-	-0.19 (0.27)	0.2 (0.23)	-0.2 (0.28)	-	-0.19 (0.27)	0.2 (0.23)	-0.2 (0.28)	-

Mediator	Behavioral intentions and behaviors											
	Change of travel plans				Get vaccinated if available				Taking protective behavior(s)			
	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test	Path a	Path b	Path c'	Sobel test
Seeking Zika information online	-0.55 (0.33)	-0.35 (0.25)	-0.03 (0.3)	-	-0.38 (0.43)	0.28 (0.32)	-0.1 (0.39)	-	-0.57 (0.33)	0.2 (0.34)	-0.48 (0.47)	-
Discussing Zika with practitioners	-1.19 (0.75)	-0.2 (0.39)	0.31 (0.27)	-	-1.05 (1.17)	0.85 (0.6)	-0.04 (0.34)	-	-1.22 (0.76)	0.78 (0.5)	-0.31 (0.38)	-
Discussing Zika with family and friends	-0.45 (0.33)	-0.06 (0.25)	0.43 (0.31)	-	-0.43 (0.45)	-0.05 (0.34)	-0.17 (0.38)	-	-0.48 (0.34)	0.84 (0.34)**	-0.95 (0.56)	-

Note. Less-educated unmarried black respondents were coded as 1 whereas other respondents were coded as -1. Unstandardized estimates and standard errors in parentheses. Main effects of sex, week number of the survey completion, seeking Zika information online, discussing Zika with practitioners, discussing Zika with family and friends, forwarding information online, reading newspapers online and offline, using online news sites, listening to radio online and offline, watching television online and offline, usage frequency of Twitter, and usage frequency of Facebook, were included for the estimations of paths *a* and *b*, while the interaction effects between the mediators and the group variable and sex were included for the estimation of paths *b* and *c'*.

*** $P \leq 0.001$. ** $P \leq 0.01$. * $P \leq 0.05$.

online and talk about Zika with practitioners, family and friends, and these factors led to higher levels of knowledge. The effects of seeking online information and talking with practitioners were stronger than the effects of discussing Zika with family and friends.

The comparison of pregnancy intentions versus current pregnancy

Table 3 shows a consistent information source differences between the pregnancy-intention- and current-pregnancy-groups. Specifically, respondents of households with members who intend to get pregnant, compared to those who are pregnant, were less likely to discuss Zika with practitioners or with family and friends (Table 3). However, seeking information online was more popular among those in the pregnancy-intention-group, $b = 0.38$, $SE = 0.19$, $P = 0.0510$. In summary, samples with pregnancy intentions (versus current pregnancy) sought information online but talked with practitioners, family and friends less. Furthermore, because information sources did not mediate differences in knowledge and behaviors, the differences in information sources had little consequence.

The unintended-pregnancy-probability-sample of younger respondents in Zika-affected regions

We also examined whether information sources differed as a function of two unintended-pregnancy-groups: Less-educated unmarried black respondents versus others. As shown in Table 4, less-educated unmarried black respondents were less likely to discuss Zika with practitioners than other respondents, $b = -1.72$ to -2.34 , $SE = 0.74-1.03$, $P = 0.0209-0.0266$. The results showed no significant associations between information sources and knowledge and behaviors measures, indicating the lack of mediating effects of information sources.

Discussion

Main finding of this study

The results from our study revealed three major findings. First, among people from households with (versus without) either current or intended pregnancy, seeking information online explained differences in behavioral intentions of changing travel plans, whereas discussing Zika with practitioners, family and friends only accounted for differences in knowledge measures (i.e. the mosquito-transmitted nature of Zika infections, the link between Zika infections and microcephaly, and the absence of noticeable symptoms in Zika infections). Second, the group with pregnancy intentions (versus current pregnancy) was less likely to discuss Zika with practitioners,

family and friends but more likely to seek information online. Third, less-educated unmarried black respondents reported fewer discussions about Zika with practitioners than other respondents, and more importantly, both groups reported inadequate knowledge about asymptomatic infections of Zika (Table 1).

What is already known on this topic

A review of the literature resulted in more than a hundred records, and about six studies examined knowledge or behaviors related to Zika in the USA and the US Virgin Islands.^{9-11,18,19} In these studies, respondents had high levels of awareness, low levels of knowledge, and varying levels of protective behaviors.^{9-11,18,19} Moreover, in prior research, people recalled hearing messages about mosquito-bites as a way of transmission but not the sexual transmission of Zika, and few recalled messages about taking specific protective behaviors.¹⁰

What this study adds

Respondents from households with members that are either currently or intending to become pregnant reported higher levels of knowledge, behavioral intentions and behaviors than those with neither current nor intended pregnancy. However, there were no differences in knowledge that Zika infection can be asymptomatic (i.e. not always having noticeable symptoms) and low levels of knowledge about sexual transmission. In line with prior studies,^{4,20} the findings are worrisome. Even though practitioners appear efficacious at increasing knowledge about Zika, as judged by lack of significant mediation, they produce very small differences in behavioral intentions and behaviors. Moreover, according to our knowledge, the present study is the first one to report differences in the use of information sources between the pregnancy-intention- and current-pregnancy-groups. Specifically, the group with pregnancy intentions was less likely to discuss Zika with practitioners, suggesting limited contacts with practitioners or inadequate access to Zika information from practitioners. More importantly, this study found differences in the unintended-pregnancy-probability-sample of younger respondents in Zika-affected regions. Less-educated unmarried black respondents, compared to others, had fewer Zika discussions with practitioners. Lastly, both groups of younger respondents reported inadequate knowledge about asymptomatic infections of Zika (Table 1).

Limitations of this study

There are two limitations in the present study. First, the survey measures of pregnancy intentions and current pregnancy

were about respondents' household members rather than the respondent themselves. Future studies may collect survey data from individual participants to determine whether our findings replicate at the individual level. Nonetheless, this study provided valuable information about households with pregnancy or intended pregnancy and their information sources. Second, the survey rotated some measures of knowledge, behavioral intentions and behaviors in and out during the period of examination, leading to the presence of systematic missing data. Future work should investigate all explanatory variables for the full period to advance existing models and theories^{21,22} in Zika infections.

Public health implications

The present study provides empirical evidence that the use of information sources differs across samples, which results in differences in knowledge, behavioral intentions and behaviors. Therefore, our findings offer a call to action for practitioners, both in the clinical and public health domains, to deliver resources to individuals at-risk for Zika infection. We recommend health campaigns to capitalize on these critical sources and thus improve knowledge about the asymptomatic nature of Zika infection and increase behavioral intentions and behaviors in vulnerable populations. These actions include encouraging practitioners to thoroughly discuss these issues, disseminating accurate information via the Internet, and opening new channels of communication for samples at risk for unintended pregnancy.

Outbreaks of Zika virus infection have been reported in Africa, the Americas, Asia and the Pacific,²³ and this widespread risk threaten women of childbearing age. More importantly, the long-term aftereffects for survivors impose high medical, societal and economic strains on the affected societies. The present findings urge all countries worldwide to make comprehensive preventive and control efforts in prenatal healthcare, family planning, and non-pregnancy related strategies. Such preventive strategies should promote discussions about Zika and the dissemination of Zika-related information in diverse channels. It is imperative to take deliberate efforts, particularly amongst pregnant women, women of childbearing age, and their partners, before an effective method is set in place to eliminate Zika vectors.

Supplementary data

Supplementary data are available at the *Journal of Public Health* online.

Conflict of interest

The study is based on a survey designed and conducted by KHJ and KW of the Annenberg Public Policy Center of the University of Pennsylvania. Two of the authors (KW and KHJ) are affiliated with the center and no authors has a conflict of interest.

Authorship

M-pSC and MF prepared the first draft of the article, conducted the literature review, and did the data analysis. KW and KHJ designed the survey on which the study is based. DA contributed to the analytic strategy and to article writing. All authors contributed in editing and revising the article.

Funding

This work was supported by the National Institutes of Health [R56 AI114501 to D.A.] and the Science of Science Communication endowment of the Annenberg Public Policy Center of the University of Pennsylvania.

References

- 1 CDC. *Zika Virus Home: Health Effects & Risks*. Centers for Disease Control and Prevention. <https://www.cdc.gov/zika/healtheffects/index.html> (17 October 2016, date last accessed).
- 2 McCarthy M. CDC updates Zika virus guidance to protect pregnant women. *Br Med J* 2016;**352**:i786.
- 3 Ellington SR, Devine O, Bertolli J *et al*. Estimating the number of pregnant women infected with Zika virus and expected infants with microcephaly following the Zika Virus outbreak in Puerto Rico, 2016. *JAMA Pediatr* 2016;**170**(10):940–5. doi:10.1001/jamapediatrics.2016.2974.
- 4 Marge Dwyer. *Many U.S. Families Considering Pregnancy don't Know Zika Facts*. Harvard T.H. Chan School of Public Health. <https://www.hsph.harvard.edu/news/press-releases/zika-virus-awareness-pregnant-women/> (19 January 2018, date last accessed).
- 5 CDC. *Areas With Zika: Case Counts in the US*. Centers for Disease Control and Prevention. <http://www.cdc.gov/zika/geo/united-states.html> (17 October 2016, date last accessed).
- 6 Meaney-Delman D, Rasmussen SA, Staples JE *et al*. Zika Virus and pregnancy: what obstetric health care providers need to know. *Obstet Gynecol* 2016;**127**(4):642–8. doi:10.1097/AOG.0000000000001378.
- 7 Vogel G. Evidence grows for Zika virus as pregnancy danger. *Science* 2016;**351**(6278):1123–4. doi:10.1126/science.351.6278.1123.
- 8 Ticconi C, Pietropoli A, Rezza G. Zika virus infection and pregnancy: what we do and do not know. *Pathog Glob Health* 2016;**110**:262–8. doi:10.1080/20477724.2016.1234804.
- 9 The Annenberg Public Policy Center. *Half of Americans Concerned Zika will Spread to Their Neighborhoods*. The Annenberg Public Policy

- Center. <http://www.annenbergpublicpolicycenter.org/half-of-americans-concerned-zika-will-spread-to-their-neighborhoods/> (19 January 2018, date last accessed).
- 10 Prue CE, Roth JN, Garcia-Williams A *et al.* Awareness, beliefs, and actions concerning Zika virus among pregnant women and community members—U.S. Virgin Islands, November–December 2016. *MMWR Morb Mortal Wkly Rep* 2017;**66**(34):909–13. doi:10.15585/mmwr.mm6634a4.
 - 11 Harvard TH Chan School of Public Health HORP. *Public Views of the Zika Virus Outbreak Topline Data*. Harvard Opinion Research Program.
 - 12 Stagg D, Hurst HM. Zika virus and pregnancy. *Nurs Womens Health* 2016;**20**(3):299–304. doi:10.1016/j.nwh.2016.03.001.
 - 13 Boulet SL, D'Angelo DV, Morrow B *et al.* Contraceptive use among nonpregnant and postpartum women at risk for unintended pregnancy, and female high school students, in the context of Zika preparedness—United States, 2011–2013 and 2015. *MMWR Morb Mortal Wkly Rep* 2016;**65**(30):780–7. doi:10.15585/mmwr.mm6530e2.
 - 14 Dehlendorf C, Rodriguez MI, Levy K *et al.* Disparities in family planning. *Am J Obstet Gynecol* 2010;**202**(3):214–20. doi:10.1016/j.ajog.2009.08.022.
 - 15 Tapales A, Finer L. Unintended pregnancy and the changing demography of American women, 1987–2008. *Demogr Res* 2015;**33**:1257–70. doi:10.4054/DemRes.2015.33.45.
 - 16 Finer LB, Zolna MR. Declines in unintended pregnancy in the United States, 2008–2011. *N Engl J Med* 2016;**374**(9):843–52. doi:10.1056/NEJMsa1506575.
 - 17 Mosher WD, Jones J, Abma JC. *Intended and Unintended Births in the United States: 1982–2010; 2012*. <http://www.ncbi.nlm.nih.gov/pubmed/23115878> (18 January 2018, date last accessed).
 - 18 Abramson D, Piltch-Loeb R. *NYU Zika Briefing Report #1. NYU Researchers Investigate Why U.S. Public Still Lacks Specific Knowledge of Zika Virus Risk*. New York City, NY; 2016. <https://www.nyu.edu/about/news-publications/news/2016/october/nyu-researchers-investigate-why-u-s-public-still-lacks-specific.html> (19 January 2018, date last accessed).
 - 19 Whittemore K, Tate A, Illescas A *et al.* Zika virus knowledge among pregnant women who were in areas with active transmission. *Emerg Infect Dis* 2017;**23**(1):164–6. doi:10.3201/eid2301.161614.
 - 20 Petersen EE, Polen KND, Meaney-Delman D *et al.* Update: Interim guidance for health care providers caring for women of reproductive age with possible Zika virus exposure—United States, 2016. *MMWR Morb Mortal Wkly Rep* 2016;**65**(12):315–322. doi:10.15585/mmwr.mm6512e2er.
 - 21 Fishbein M, Cappella JN. The role of theory in developing effective health communications. *J Commun* 2006;**56**(SUPPL). 10.1111/j.1460-2466.2006.00280.x.
 - 22 Petty RE, Cacioppo JT. The elaboration likelihood model of persuasion. In: Berkowitz L (ed). *Advances in Experimental Social Psychology*; Vol 19. Orlando, FL: Academic Press, 1986:123–205. doi:10.1016/S0065-2601(08)60214-2.
 - 23 WHO. *Zika Virus Infection: Disease Outbreak News*. World Health Organization. <http://www.who.int/csr/don/archive/disease/zika-virus-infection/en/> (14 April 2018, date last accessed).